

ORAL PRESENTATION

Open Access

O005: Results of the french national audit on standard precautions

M Giard^{1,2*}, E Laprugne-Garcia¹, E Caillat-Vallet¹, I Russell¹, D Verjat-Trannoy³, M-A Ertzscheid⁴, N Vernier⁵, C Laland⁶, A Savey^{1,2}, GREPHH¹

From 2nd International Conference on Prevention and Infection Control (ICPIC 2013) Geneva, Switzerland. 25-28 June 2013

Introduction

Standard precautions (SP) aim to protect healthcare workers (HCW) and patients from infectious diseases arising from bloodborne pathogens and reduce the risk of cross transmission of micro-organisms. They must be applied in all circumstances, regardless of the infectious status of the patient.

Objectives

The objectives were to assess: 1) institutional policies for SP promotion; 2) available resources for SP implementation; and 3) education of HCW and their compliance with SP.

Methods

The study was a mixed audit of procedures, resources and attitudes. It was conducted between February 1st and December 31st 2011, supported by the Ministry of Health. Inclusion criteria were voluntary public and private hospitals in France, medical, surgical and medico-technical wards therein and HCW working with patients in these wards. Self-assessment questionnaires were administered at three levels: institutional, ward and HCW. At institutional and ward levels, results were given as a percentage of objectives attained; at professional level, percentages of responses reported as “never”, “sometimes”, “often” or “always” were calculated for each question.

Results

A total of 1,599 hospitals participated, including 14,968 wards and 203,840 HCW. At institutional level, the overall score was 88%, covering: SP promotion (91%),

procedures (99%) and SP evaluation (63%). At ward level, the overall score was 94%, covering: procedures (95%) and resources (93%). Among the 165,722 (81.3%) HCW who reported having participated in a training session on SP, 69.6% had had it in the last five years. A total of 88.1% of HCW knew where to find the appropriate written procedure in the event of a blood exposure. HCW reported the best compliance for glove changing between two patients (94.5% “always”). The less respected criteria were glove use for intramuscular or subcutaneous injection and eye protection use in the event of blood exposure risk (34.5% and 24.4% “always”, respectively).

Conclusion

No study on SP exists in literature which includes such a large participation as this one. It will form a reference basis leading to actions for improvement at local and national level.

Disclosure of interest

None declared.

Author details

¹CClin Sud-Est, Lyon, France. ²UMR5558, LBBE, Lyon 1 University, Villeurbanne, France. ³CClin Paris-Nord, Paris, France. ⁴CClin Ouest, Rennes, France. ⁵CClin Est, Nancy, France. ⁶CClin Sud-Ouest, Poitiers, France.

Published: 20 June 2013

doi:10.1186/2047-2994-2-S1-O5

Cite this article as: Giard et al.: O005: Results of the french national audit on standard precautions. *Antimicrobial Resistance and Infection Control* 2013 **2**(Suppl 1):O5.

¹CClin Sud-Est, Lyon, France

Full list of author information is available at the end of the article