

POSTER PRESENTATION

Open Access

P323: Epidemiological study of drug administration routes (dar) in the department of pediatrics, Gabriel Touré hospital. Mali

M Daouda^{1*}, AS Kaya², B Traoré³, L Bengaly⁴, M Sacko⁵, A Camara¹, AM Traoré¹, M Diakit¹, H Cissé¹, AT Sidibé², HA Traoré², T Sidibé³, MM Keita³

From 2nd International Conference on Prevention and Infection Control (ICPIC 2013) Geneva, Switzerland. 25-28 June 2013

Objectives

Injectable route seems to be the most frequently used in health facilities in a resource-limited setting.

Methods

In order to better understand this under investigated issue, we conducted a longitudinal study of DAR use in the general pediatric ward of the Gabriel Touré Hospital to Bamako during 6 months.

Results

We are interested in the routes of administration applied to a population of 300 children with a sex ratio (M/F) = 1.3. Their average age was 2 years \pm 1. Presumed diagnoses underwent a change from admission to discharge, both in frequency and formulation. Malaria (37.4% vs 39.7% at 72 hours), pneumonia (19% vs 20% at 72 hours), and the nephrotic syndrome (2.2% vs. 5.1% in 72 hours) were most commonly mentioned. Treatments prescribed for the presumed diagnosis were administered parenterally in 76.6% of cases at admission, in 70% 72h hours after admission and in 36.3% at discharge. The following complications were noted: inflammation of the catheter puncture sites (21.8% at admission, 18% after 5 days of hospitalization), abscess at the site of intramuscular injection (2.1%). The mean duration of hospitalization was 7.6 \pm 3.7 days and mortality was 11%.

Conclusion

Injection is the most widely used in the pediatric unit III for the most common pathologies. A detailed study

would be needed to assess the adequacy of the diagnostic hypotheses and routes of administration. The high number of injections exposes staff and patients to risks.

Disclosure of interest

None declared.

Author details

¹Service of Infectious Diseases, Department of Public Health, Faculty of Medicine, Bamako, Mali. ²Department of Internal medicine, University Hospital of Point G, Department of Public Health, Faculty of Medicine, Bamako, Mali. ³Pediatric Services, Department of Public Health, Faculty of Medicine, Bamako, Mali. ⁴Hospital Pharmacy, CHU Gabriel Touré, Department of Public Health, Faculty of Medicine, Bamako, Mali. ⁵Department of Public Health, Faculty of Medicine, Bamako, Mali.

Published: 20 June 2013

doi:10.1186/2047-2994-2-S1-P323

Cite this article as: Daouda et al.: P323: Epidemiological study of drug administration routes (dar) in the department of pediatrics, Gabriel Touré hospital. Mali. *Antimicrobial Resistance and Infection Control* 2013 **2**(Suppl 1):P323.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

