

POSTER PRESENTATION

Open Access

Prevalence of colonisation with vancomycin-resistant enterococci on admission - a cross-sectional study in 6 German university hospitals, 2014

M Wiese-Posselt^{1*}, J Zweifgner², AM Rohde³, F Schwab³, W Kern⁴, H Seifert², P Gastmeier³,
the DZIF-ATHOS Study Group

From 3rd International Conference on Prevention and Infection Control (ICPIC 2015)
Geneva, Switzerland. 16-19 June 2015

Introduction

The survey presented here is part of the multicenter study ATHOS (antibiotic therapy optimisation study) on prevalence and incidence of nosocomial carriage of multi-drug resistant organisms (MDROs) in Germany.

Objectives

The aim of this survey was to assess the prevalence of rectal colonisation with VRE in patients on hospital admission. Additionally, we performed a risk factor analysis for VRE carriage.

Methods

We recruited adult patients within 72 h after admission to non-intensive care units in six German university hospitals to obtain rectal swabs that were screened for VRE. Each patient was asked to answer a short questionnaire on potential risk factors for colonisation with MDROs, such as sex, age, antibiotic therapy in the last 6 months, previous colonisation with any MDRO, travel abroad in the last 6 months, stay in a nursing home or hospitalisation in the last 6 months. Univariable and multivariable risk factor analyses were performed to identify those factors which were associated with VRE colonisation.

Results

In 2014, 4372 patients were included in the admission survey. Overall, 35 patients were colonised with VRE (admission prevalence of 0.8%). The following factors were

associated significantly with VRE carriage: centre (OR = 3.61, CI 95% 3.05-4.27, $p < 0.001$), previous MDRO colonisation (OR = 3.23, CI 95% 1.47-7.11, $p = 0.004$), antibiotic use (OR = 2.91, CI 95% 1.47-5.74, $p = 0.002$), stay in a nursing home (OR = 3.19, CI 95% 1.87-5.46, $p < 0.001$), and hospitalisation (OR = 3.19, CI 95% 2.06-5.42, $p = 0.000$). Interestingly, travel abroad had a protective effect on VRE carriage (OR = 0.51, CI 95% 0.29-0.91, $p < 0.001$).

Conclusion

This admission survey in Germany revealed a moderate colonisation rate of 0.8%, whereat prevalence differed significantly by centre. Well-known risk factors for VRE carriage were identified: previous colonisation with any MDRO, antibiotic use in the last 6 months, and admission to nursing homes or hospitals in the last 6 months. We observed that not travelling abroad was significantly associated with VRE carriage which we cannot explain. Thus, we cannot exclude a confounding factor we have not considered so far and should be addressed in further studies.

Disclosure of interest

None declared.

Authors' details

¹Institute of Hospital Hygiene, Charité, Berlin, Germany. ²University Hospital Cologne, Germany. ³Charité, Germany. ⁴University Hospital Freiburg, Germany.

¹Institute of Hospital Hygiene, Charité, Berlin, Germany
Full list of author information is available at the end of the article

Published: 16 June 2015

doi:10.1186/2047-2994-4-S1-P198

Cite this article as: Wiese-Posselt *et al.*: Prevalence of colonisation with vancomycin-resistant enterococci on admission - a cross-sectional study in 6 German university hospitals, 2014. *Antimicrobial Resistance and Infection Control* 2015 **4**(Suppl 1):P198.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

