


SHORT REPORT

Open Access



Preparing for future waves and pandemics: a global hospital survey on infection control measures and infection rates in COVID-19

Simon Matoori^{1,2,3*} , Daniel R. Kuritzkes⁴, Yonggeng Goh⁵, Swee Tian Quek⁵, Liang Wang⁶, Ziyun Sun⁶, Fabiano di Marco^{7,8}, Daniela Borleri⁹, Sabrina Buoro⁹, Stefano Fagioli¹⁰, Tatiana Ferrari¹¹, Marco Rizzi¹², Federico Raimondi¹³, Simonetta Cesa¹⁴, Dow-Mu Koh³, Johannes M. Froehlich¹⁵, Sonja Janssen¹⁶, Bettina Lange¹⁷, Alexander Egle¹⁸, Stefan Erb¹⁹, Erik Mossdorf¹⁹ and Andreas Gutzeit^{2*}

Abstract

A survey of hospitals on three continents was performed to assess their infection control preparedness and measures, and their infection rate in hospital health care workers during the COVID-19 pandemic. All surveyed hospitals used similar PPE but differences in preparedness, PPE shortages, and infection rates were reported.

Introduction

In the COVID-19 pandemic, the need for personal protective equipment (PPE) strongly increased to protect health care workers (HCWs) within the hospital from infection by COVID-19-positive patients [1]. Although our understanding of the novel coronavirus strain SARS-CoV-2 has been rapidly evolving, a detailed understanding on the modes of transmission is still lacking [2, 3]. To protect HCWs, the World Health Organization and the Centers for Disease Control recommend the use of face masks, eye protection, gowns, medical gloves, frequent hand hygiene, environmental cleaning, and waste management [4]. The most potent protection, respirators which filter out at least 94% (FFP2, N95) or 99% (FFP3, N99) of airborne particles, as well as other PPE were quickly in short demand as the pandemic swept around the planet, resulting in shortages in many hospitals with

a potential increase in COVID-19 infection risk for HCWs [5, 6].

Our hypothesis was that the preparedness and response to the pandemic was different among hospitals, countries, and regions, and resulted in different exposure risk for HCWs in terms of COVID-19 prevalence and PPE availability. The goal of our study was to compare and assess the preparedness, the response, and the infection rate among hospital HCWs of several institutions on three continents by questionnaire.

Methods

No Institutional Review Board approval or informed patient consent was needed for this survey. The survey was performed between June and August 2020 at eight different institutions: Salzburg (Austria), Wuhan (China), Mannheim (Germany), Bergamo (Italy), Singapore, Hirslanden Klinik in Lucerne (Switzerland), an anonymized institution in the UK, Boston (MA, USA). Sites were recruited by emailing the Division of Infectious Diseases of each hospital. Surveys were sent out by email and reports were received by email.

*Correspondence: simon.matoori@umontreal.ca; andreas.gutzeit@hirslanden.ch

¹ Faculté de Pharmacie, Université de Montréal, 2940 Chemin de Polytechnique, Montréal, QC H3T 1J4, Canada

² Department of Health Sciences and Medicine, University of Lucerne, Lucerne, Switzerland

Full list of author information is available at the end of the article



Questionnaire

The surveyed hospitals were given a questionnaire about institutional baseline characteristics and infection control measures for the COVID-19 pandemic of 2020. The questionnaire was answered by an infectious disease expert of the corresponding institution, and included the following topics: (1) Baseline characteristics of the participating hospitals; (2) COVID-19 infection rates among HCWs, (3) Pandemic preparations and trainings for a pandemic with an infectious respiratory disease before the COVID-19 outbreak, (4) Personal protective equipment (PPE) for health care worker with close contact with COVID-19 positive or suspected patients during the COVID-19 pandemic, and (5) various management issues of the COVID-19 pandemic (Additional file 1).

The infection rate was calculated as the number of infected HCWs divided by the total number of HCWs in the hospital. Shortages in PPE were defined as the absence of the PPE article in question. Survey respondents retrieved all hospital infection data reported in this study from their hospital records.

Results

Baseline characteristics of the surveyed hospitals and COVID-19 infection rates

The overall survey response rate was 100%. No incomplete survey responses were received.

Baseline characteristics of the eight hospitals included in the survey and the COVID-19 case load of their country are shown in Table 1 and Additional file 1: Table S1. Five hospitals are located in Europe (Austria, Germany, Italy, Switzerland, United Kingdom), two in Asia (China, Singapore) and one in North America (United States) (Additional file 1: Fig. S1).

All surveyed hospitals are mid-sized to large tertiary referral centers. The percentage of COVID-19-positive

HCWs varied widely from 0.1 to 8.5%. The two Asian institutions showed low infection rates ($\leq 1\%$) despite treating large numbers of COVID-19 positive patients.

Pandemic preparedness before the COVID-19 outbreak

All hospitals except two (Italy and United Kingdom) reported respiratory infectious disease outbreak preparedness with basic infection control trainings of any kind before the onset of the COVID-19 pandemic (Additional file 1: Table S2).

The institutions in China and Singapore noted that their plan for a pandemic with a respiratory infection was mainly based on the experience from the first SARS epidemic and the resulting policies. While the institution in Italy did not specifically prepare for a pandemic with a respiratory infectious disease, there was a course to prepare for an Ebola outbreak aimed at the Emergency Department in 2019. Furthermore, there was an emergency preparation meeting before the first COVID-19 patient was admitted to the Italian institution.

Personal protective equipment during the COVID-19 outbreak in case of close contact with COVID-19 patients

All surveyed hospitals reported the use of surgical masks or respirators (FFP2/3, N95) and other disposable PPE such as gloves, gowns, and hand disinfectants (Table 2). Disposable caps and goggles were not used in two European hospitals. Other PPE used include face shields, visors, aprons, and air-purifying respirators in aerosolizing procedures. HCWs were trained to put on and use the PPE in all institutions.

Four of the eight surveyed institutions reported shortages in PPE at some point in the pandemic (Additional file 1: Table S3). Masks or respirators were missing for at least a brief period of time at the institutions in Austria, Germany, Italy, and the US. Three of the eight institutions

Table 1 Hospital baseline information and number of infected HCWs and patients

Institution	Total hospital beds	Total HCWs	Total infected HCWs	Percentage of infected HCWs (%)	Number of treated COVID-19 patients	Date of questionnaire completion
Salzburg, Austria	1167	3471	43	1.2	170	June 5, 2020
Wuhan, China	7000	9700	100	1.0	3200	July 23, 2020
Mannheim, Germany	1300	3719	17	0.4	63	June 3, 2020
Bergamo, Italy	1000	4705	398	8.5	2141 ^a	September 24, 2020
Singapore	1160	7800	7	0.1	887	June 2, 2020
Lucerne, Switzerland	220	1300	6	0.4	18	August 10, 2020
UK	269	3900	278	7.1	95	August 1, 2020
Boston, USA	793	24,000	450	1.9	682 ^b	August 18, 2020

^a Number of hospital-admitted COVID-19 patients (total number in Emergency Department: 4212)

^b Until June 30, 2020

Table 2 Availability of PPE for frontline HCWs during the COVID-19 pandemic

Institution	Masks		Disposable gloves	Disposable gowns	Disposable caps	Disposable goggles	Hand disinfectant	Other PPE	Training
	Type	Duration							
Salzburg, Austria	SM, FFP2, FFP3	4 h	Yes	Yes	Yes	Yes ^a	Yes	Face shields	Yes
Wuhan, China	SM, N95, KN95	4 h	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Mannheim, Germany	FFP2, FFP3	1/shift	Yes	Yes	Yes	Yes	Yes	Aprons	Yes
Bergamo, Italy	SM, FFP2, FFP3, N95, N99, KN95	1–2/shift	Yes	Yes	Yes	Yes	Yes	Face shields	Yes
Singapore	SM, N95	4–5 h	Yes	Yes	Yes	Yes	Yes	Powered air-purifying respirator (interventional radiology)	Yes
Lucerne, Switzerland	SM, FFP2	1/shift	Yes	Yes	No	Yes	Yes	No	Yes ^b
UK	SM, FFP2, FFP3	1/shift	Yes	Yes	Yes	No	Yes	Face shields	Yes ^c
Boston, USA	N95 (fitted)	1/shift (8–12 h)	Yes	Yes	No	Yes ^a	Yes	Personal respirators (if unable to use N95), face shields ^a	Yes

SM surgical masks

^a Reused after disinfection

^b Only HCWs in risk areas (COVID-19 ward, emergency department, intensive care unit)

^c Only frontline staff with contact with COVID-19 patients

reported that disposable gowns were not available during the entire time of the pandemic. Single institutions further reported shortages in gloves and disinfectants, and the need for rationing PPE to avoid shortages.

Management of the pandemic

Infectious disease specialists were in charge of the management of the pandemic in all surveyed hospitals except for the one in the UK where non-specialized physicians were responsible (Additional file 1: Table S4). All hospitals put a task force in charge of coordinating the safety measures to reduce viral spread in the hospital, and implemented zoning (i.e., areas designated for suspected and confirmed COVID-19 patients), grouping COVID-19-positive patients (cohorting), hygiene trainings, and restrictions for visitors. Masking of HCWs in the entire hospital was mandated by all hospitals except for the one in Salzburg, where it was recommended. Visitor restrictions were put in place in all surveyed hospitals, with two hospitals allowing no visitors and two only allowing compassionate visits.

Discussion

In this study, we evaluated the preparedness, the management, and the infection rate of hospital HCWs of the COVID-19 pandemic of eight hospitals located in Europe, Asia, and North America. While most hospitals

were prepared for a pandemic with a respiratory disease, there were shortages in several PPE, namely masks and gowns. The percentage of COVID-19-positive hospital workers ranged from 0.1 to 8.5%. Notably, the Asian hospitals reported no shortages in surgical masks or respirators and showed low infection rates ($\leq 1\%$) despite high numbers of treated COVID-19 patients. Without pointing fingers, this survey helps us identify key aspects of preparedness in terms of planification and PPE item stockage, in order to be better prepared for future waves of COVID-19 and other pandemics with respiratory viruses.

The low infection numbers despite high caseloads in Asian hospital may be explained by the experience with the SARS epidemic in 2002/2003 which mainly affected Asian countries such as China, Singapore, Vietnam, and Hong Kong. Hospital transmission accounted for a 22 to 57% of SARS cases [7], and was primarily related to inappropriate or missing PPE, hand disinfection, and hospital personnel education (e.g., removal of contaminated PPE) [8]. As a result, these countries prepared sophisticated plans to manage future pandemics, mandating training hospital personnel for pandemics with respiratory pathogens, stockpiling PPE, and further measures such as zoning [9]. As no shortages in masks and respirators were reported by these hospitals, their stockpiling has likely contributed to the low infection

rates in these hospitals, which underlines their importance in protecting HCWs from COVID-19 patients.

In our survey, countries with less exposure to the first SARS epidemic, namely in Europe and the US, were less prepared in terms of personnel training and/or PPE stockpiling, and experienced more PPE shortages and higher numbers of infected hospital HCWs. However, hospitals showed a steep learning curve, adopting measures such as universal masking for hospital personnel and proving their usefulness for infection control [10–12].

As we prepare for the next waves of COVID-19, we have learned that universal masking of hospital personnel with surgical masks, immediately isolating COVID-19-positive patients, and the use of highly protective PPE (respirators, eye protection, gowns) in contact with COVID-19 patients is of high utility to protect HCWs [4, 13]. As some of these insights were already available after the SARS epidemic and improved the response of the surveyed Asian hospitals to the COVID-19 pandemic, it is instrumental for hospitals to learn from each other's experiences and to cooperate to improve hospital infection control in future waves of COVID-19 and other pandemics. Based on the information of this survey, it will be essential to either stock higher amounts of PPE or to have the capacity to produce or recycle PPE locally to be prepared for future disruptions in supply chains.

This study has several limitations. Most importantly, we cannot determine if the COVID-19 positive hospital HCWs were infected in the hospital or outside. Indeed, the true extent of COVID-19 is difficult to assess [14, 15]. Another limitation of the study is that the questionnaire was filled out when some countries were still in the first wave while others were past the first wave (e.g., China). Furthermore, there might be a selection bias as most hospitals were selected based on previous collaborations. Another limitation is that all surveyed hospitals are located in upper-middle to high-income countries. Finally, some surveyed European hospitals treated low numbers of COVID-19 patients, complicating the evaluation of their true preparedness.

In this survey, all responding hospitals in Asia, Europe, and North America reported the use of recommended PPE but the degree of preparedness, PPE shortages, and infection rate among hospital HCWs varied widely. Our study highlights the importance of universal masking of HCWs with surgical masks, immediately isolating COVID-19-positive patients, and the use of highly protective PPE (respirators, eye protection, gowns). To prepare for the next waves of COVID-19 and future pandemics, we further urge

hospitals to learn from each other and to cooperate to improve infection control.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13756-021-01029-z>.

Additional file 1: Supplementary Figure and Tables.

Acknowledgements

We thank Health Care Workers worldwide for their efforts during the COVID-19 pandemic.

Authors' contributions

SM, SE, and AG prepared the questionnaire and wrote the original draft of the manuscript. All authors contributed to the data presented in this manuscript, edited the manuscript, and approved the final version of the manuscript.

Funding

All authors report no financial support relevant to this manuscript.

Availability of data and materials

All relevant data and material are included in the main text and the supporting information of this manuscript.

Declarations

Ethics approval and consent to participate

Ethics approval was waived for this survey.

Consent for publication

All authors consented to the publication of this manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Faculté de Pharmacie, Université de Montréal, 2940 Chemin de Polytechnique, Montreal, QC H3T 1J4, Canada. ²Department of Health Sciences and Medicine, University of Lucerne, Lucerne, Switzerland. ³Department of Radiology, Paracelsus Medical University, Salzburg, Austria. ⁴Division of Infectious Diseases, Brigham and Women's Hospital, Boston, MA, USA. ⁵Department of Diagnostic Imaging, National University Hospital, Singapore, Republic of Singapore. ⁶Department of Radiology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China. ⁷Pneumology Unit, Papa Giovanni XXIII Hospital, Bergamo, Italy. ⁸Health Sciences Department, University of Milan, Milan, Italy. ⁹Occupational Medicine Unit, Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹⁰Quality Management Unit, Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹¹Gastroenterology, Hepatology and Liver Transplantation, Department of Medicine, Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹²Infectious Diseases Unit, Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹³Respiratory Unit, Department of Medicine, Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹⁴Department of Health and Social Care Professions, Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹⁵Clinical Research Group, Klus Apotheke Zurich, Zurich, Switzerland. ¹⁶Clinic of Radiology and Nuclear Medicine, University Medical Center Mannheim, University of Heidelberg, Mannheim, Germany. ¹⁷Hospital Hygiene Unit, University Medical Center Mannheim, University of Heidelberg, Mannheim, Germany. ¹⁸Ilrd Medical Department with Hematology and Medical Oncology, Hemostaseology, Rheumatology and Infectious Diseases, Oncologic Center, Paracelsus Medical University, Salzburg, Austria. ¹⁹Department of Infectious Diseases, Hirslanden Klinik St. Anna, Lucerne, Switzerland.

Received: 12 October 2021 Accepted: 24 October 2021

Published online: 20 December 2021

References

1. Kameron D. Covid-19: the crisis of personal protective equipment in the US. *BMJ*. 2020;369:1367.
2. Meselson M. Droplets and aerosols in the transmission of SARS-CoV-2. *N Engl J Med*. 2020;382(21):2063.
3. Fennelly KP. Particle sizes of infectious aerosols: implications for infection control. *Lancet Respir Med*. 2020;8(9):914–24.
4. Bartoszko JJ, Farooqi MAM, Alhazzani W, Loeb M. Medical masks vs N95 respirators for preventing COVID-19 in healthcare workers: a systematic review and meta-analysis of randomized trials. *Influenza Other Respi Viruses*. 2020;14(4):365–73.
5. Ranney ML, Griffith V, Jha AK. Critical supply shortages—the need for ventilators and personal protective equipment during the Covid-19 pandemic. *N Engl J Med*. 2020;382(18):E41.
6. Livingston E, Desai A, Berkwitz M. Sourcing personal protective equipment during the COVID-19 pandemic. *JAMA*. 2020;323(19):1912–4.
7. Chowell G, Abdirizak F, Lee S, et al. Transmission characteristics of MERS and SARS in the healthcare setting: a comparative study. *BMC Med*. 2015;13(1):210.
8. Shaw K. The 2003 SARS outbreak and its impact on infection control practices. *Public Health*. 2006;120(1):8–14.
9. Goh Y, Chua W, Lee JKT, et al. Operational strategies to prevent coronavirus disease 2019 (COVID-19) spread in radiology: experience from a Singapore Radiology Department after severe acute respiratory syndrome. *J Am Coll Radiol*. 2020;17(6):717–23.
10. Wang X, Ferro EG, Zhou G, Hashimoto D, Bhatt DL. Association between universal masking in a health care system and SARS-CoV-2 positivity among health care workers. *JAMA*. 2020;324(7):703.
11. Klompas M, Morris CA, Sinclair J, Pearson M, Shenoy ES. Universal masking in hospitals in the COVID-19 era. *N Engl J Med*. 2020;382(21):e63.
12. Gutzeit A, Li Q, Matoori S, Li B, Wang L. What can European radiologists learn from the outbreak of COVID-19 in China? A discussion with a radiologist from Wuhan. *Eur Radiol*. 2020;30(7):3609–11.
13. Qaseem A, Etcheandia-Ikobaltzeta I, Yost J, et al. Use of N95, surgical, and cloth masks to prevent COVID-19 in health care and community settings: living practice points from the American College of Physicians (version 1). *Ann Intern Med*. 2020;178:M20-3234.
14. Matoori S, Khurana B, Balcom MC, Koh DM, Froehlich JM, Janssen S, Kolokythas O, Gutzeit A. Intimate partner violence crisis in the COVID-19 pandemic: how can radiologists make a difference? *Eur Radiol*. 2020;30(12):6933–6.
15. Matoori S, Khurana B, Balcom MC, Froehlich JM, Janssen S, Forstner R, King AD, Koh DM, Gutzeit A. Addressing intimate partner violence during the COVID-19 pandemic and beyond: how radiologists can make a difference. *Eur Radiol*. 2021;31(4):2126–31.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

